



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

FREE PURE CULTURES OF AMEBÆ PARASITIC IN MAMMALS

Williams (Jour. Med. Research, Dec., 1911) gives a summary of the results of efforts to get pure cultures of amebæ found in human and other mammalian bodies, and to cultivate these without the bacteria which accompany them.

1. Certain strains of such parasitic amebæ may readily be kept going by using as food sterilized brain, liver, or kidney tissue from dogs, rabbits or guinea-pigs.
2. Such artificial cultures do not show essential morphologic variations as the result of this change of regimen.
3. Certain characteristics used by some students as diagnostic of the pathogenic forms (as, large size, motility, highly refractive ectoplasm, weak, chromatin, etc.) are found to belong as well to non-pathogenic forms, apparently; and thus the diagnostic value of these qualities is correspondingly diminished.
4. It is suggested that the amebæ themselves, and not merely the bacteria usually associated with them, may be responsible for amebic dysentery and amebic abscesses.

A METHOD OF DIAGNOSIS IN SYPHILIS

Noguchi (Jour. Exp. Med., Dec., 1911) reports instances of a cutaneous reaction produced in syphilitic patients by the injection of *luctin* (an extract of pure cultures of *Treponema pallidum*), especially in tertiary and hereditary (latent) cases, which he hopes may prove valuable in diagnosis of this disease,—as supplementing the Wassermann reaction which is more constant in primary or secondary forms.

TREPONEMA PALLIDUM IN MONKEYS

The same investigator (Jour. Exp. Med., Jan., 1912) has inoculated two species of lower monkeys with pure cultures of *T. pallidum* taken originally from human syphilitic lesions. This results in lesions resembling the primary lesions in man and those caused in the monkey by direct inoculation of monkeys with human serum containing the spirochaetes. The blood of the monkeys furthermore developed the property of giving the positive Wassermann reaction. This result verifies the identity of his supposed pure strains with the disease germ, and definitely relates the Wassermann reaction to the *T. pallidum*.